

# Math 213 - Maximum and Minimum Values, II

Peter A. Perry

University of Kentucky

October 10, 2018

# Homework

- Re-read section 14.7, esp. pp. 965–967
- Continue homework on section 14.7, problems 1-15 (odd), 31, 33, 37, 41-49 (odd)
- Read section 14.8 for Friday's lecture
- Finish webwork B4 on 15.5-14.6 (chain rule, directional derivatives)
- Study for Quiz #6 on 14.5-14,7
- Remember that Exam II is next Wednesday, October 17 at 5 PM

## Unit II: Differential Calculus of Several Variables

- Lecture 13 Functions of Several Variables
- Lecture 14 Limits and Continuity
- Lecture 15 Partial Derivatives
- Lecture 16 Tangent Planes and Linear Approximation, I
- Lecture 17 Tangent Planes and Linear Approximation, II
- Lecture 18 The Chain Rule
- Lecture 19 Directional Derivatives and the Gradient
- Lecture 20 Maximum and Minimum Values, I
- Lecture 21 **Maximum and Minimum Values, II**
- Lecture 22 Lagrange Multipliers
- Lecture 23 Review for Exam 2

# Goals of the Day

- Understand how to find absolute maxima and minima of functions of two variables on a bounded, closed set

# Review of Calculus I

**The Closed Interval Method** To find the *absolute* maximum and minimum values of a continuous function on a closed interval  $[a, b]$ :

1. Find the values of  $f$  at the critical numbers of  $f$  in  $[a, b]$
2. Find the values of  $f$  at the endpoints of the interval
3. The largest of the values from steps 1 and 2 is the absolute maximum of  $f$  on  $[a, b]$ ; the smallest of these values is the absolute minimum of  $f$  on  $[a, b]$ .


For functions of two variables:

1. The “closed interval” on the line is replaced by a “closed set” in the plane
2. The boundary of a closed set is a *curve* rather than just two points


Otherwise, the idea is much the same!

# Bounded Sets, Closed Sets, Boundaries

A *bounded set*  $D$  in  $\mathbb{R}^2$  is a set that can be enclosed inside a large enough circle




# Bounded Sets, Closed Sets, Boundaries



A *bounded set*  $D$  in  $\mathbb{R}^2$  is a set that can be enclosed inside a large enough circle

A *boundary point* is a point  $(a, b)$  that belongs to  $D$  but has points that don't belong to  $D$  arbitrarily close to it

# Bounded Sets, Closed Sets, Boundaries




A *bounded set*  $D$  in  $\mathbb{R}^2$  is a set that can be enclosed inside a large enough circle

A *boundary point* is a point  $(a, b)$  that belongs to  $D$  but has points that don't belong to  $D$  arbitrarily close to it

The *boundary* of a set  $D$  is the set consisting of all the boundary points



# Bounded Sets, Closed Sets, Boundaries



A *bounded set*  $D$  in  $\mathbb{R}^2$  is a set that can be enclosed inside a large enough circle

A *boundary point* is a point  $(a, b)$  that belongs to  $D$  but has points that don't belong to  $D$  arbitrarily close to it

The *boundary* of a set  $D$  is the set consisting of all the boundary points

A *closed set*  $D$  is one that contains all of its boundary points.

# Bounded Sets, Closed Sets, Boundaries

Classify each of the following sets as bounded or not bounded, and closed or not closed

1.  $D = \{(x, y) : x^2 + y^2 < 1\}$
2.  $D = \{(x, y) : x^2 + y^2 \leq 1\}$
3.  $D = \{(x, y) : x^2 + y^2 \geq 1\}$
4.  $D = \{(x, y) : x^2 + y^2 > 1\}$

# The Extreme Value Theorem

**Extreme Value Theorem** If  $f$  is continuous on a closed, bounded set  $D$  in  $\mathbb{R}^2$ , then  $f$  attains an absolute maximum value  $f(x_1, y_1)$  and an absolute minimum value  $f(x_2, y_2)$  at some points  $(x_1, y_1)$  and  $(x_2, y_2)$  in  $D$ .

*Practical fact:* These extreme values occur either in the interior of  $D$ , where the second derivative test works, or on the boundary of  $D$ , where the search for maxima and minima can be reduced to a Calculus I problem.


# The Closed Set Method

**The Closed Set Method** To find the absolute minimum and maximum values of a continuous function  $f$  on a closed, bounded set  $D$ :

1. Find the values of  $f$  at critical points of  $f$  in  $D$
2. Find the extreme values of  $f$  on the boundary of  $D$
3. The largest of the values from steps 1 and 2 is the absolute maximum value; the smallest of these values is the absolute minimum value.

The tricky bit is step 2.


## Warm-Up: Finding Extreme Values on a Boundary



1. Find the extreme values of

$$f(x, y) = x^2 - y^2$$

on the boundary of the disc  
 $x^2 + y^2 = 1$




2. Find the extreme values of

$$f(x, y) = x^2 + y^2 - 2x$$

on the boundary of the rectangular  
region with vertices  $(2, 0)$ ,  $(0, 2)$  and  
 $(0, -2)$ .

# Finding Extreme Values




1. Find the extreme values of

$$f(x, y) = x^2 - y^2$$

on the disc

$$\{(x, y) : x^2 + y^2 \leq 1\}.$$




2. Find the extreme values of

$$f(x, y) = x^2 + y^2 - 2x$$

on the rectangular region with vertices  $(2, 0)$ ,  $(0, 2)$  and  $(0, -2)$ .

## More Extreme Values




Find the absolute maximum and absolute minimum of

$$f(x, y) = xy^2$$

on the region

$$D = \{(x, y) : x \geq 0, y \geq 0, x^2 + y^2 \leq 3\}$$

# Yet More Extreme Values




Find the absolute maximum and absolute minimum of

$$f(x, y) = x^3 - 3x - y^3 + 12y$$

if  $D$  is the quadrilateral whose vertices are  $(-2, 3)$ ,  $(2, 3)$ ,  $(2, 2)$ , and  $(-2, -2)$ .



# A Word Problem with Extreme Values



Find the volume of the largest rectangular box in the first octant with three faces in the coordinate planes and one vertex in the plane

$$x + 2y + 3z = 6.$$

1. What is the volume of the box in terms of  $(x, y)$  only?
2. What values of  $(x, y)$  are allowed?
3. Do we need to check the boundary?