▲□▶ ▲圖▶ ★ 国▶ ★ 国▶ - 国 - のへで

Math 213 - Tangent Planes and Linear Approximation

Peter A. Perry

University of Kentucky

February 13, 2019



Homework

- Remember that WebWorks B1 and B2 are due tonight! (Pay special attention to problem 7)
- Re-read section 14.4
- Start working on practice problems in section 14.4, 1, 3, 5, 11-21 (odd), 25-33 (odd)
- Prepare for quiz on sections 14.1 and 14.3 tomorrow
- Read section 14.5

▲ロト ▲帰 ト ▲ ヨ ト ▲ ヨ ト ・ ヨ ・ の Q ()

Unit II: Differential Calculus of Several Variables

- Lecture 12 Functions of Several Variables
- Lecture 13 Partial Derivatives
- Lecture 14 Tangent Planes and Linear Approximation
- Lecture 15 The Chain Rule
- Lecture 16 Directional Derivatives and the Gradient
- Lecture 17 Maximum and Minimum Values, I
- Lecture 18 Maximum and Minimum Values, II
- Lecture 19 Lagrange Multipliers
- Lecture 20 Double Integrals
- Lecture 21 Double Integrals over General Regions
- Lecture 22 Double Integrals in Polar Coordinates

Lecture 23 Exam II Review

Goals of the Day

- Understand how the partial derivatives $f_x(a, b)$ and $f_y(a, b)$ define the *tangent plane* to the graph of z = f(x, y) at (a, b, f(a, b))
- Understand how the partial derivatives f_x(a, b) and f_y(a, b) define the *linear approximation* L(x, y) to f(x, y) near (x, y) = (a, b)
- Understand the *total differential dz* of a function z = f(x, y) and how it's used to compute percentage change and analyze error
- Generalize these ideas to functions of three variables

Warm-Up: Linear Functions



The graph of a line Ax + By = C defines a *linear function* of one variable

$$y = f(x) = \frac{C}{B} - \frac{A}{C}x$$

The graph of a plane ax + by + cz = d defines a *linear function* of two variables

$$z = f(x, y) = \frac{d}{c} - \frac{a}{c}x - \frac{b}{c}y$$

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ ○臣 - の々ぐ

Functions of One Variable - Tangent Line



The derivative f'(a) gives the slope of the tangent line to the graph of y = f(x) at (a, f(a)).

The derivative f'(a) defines a linear function

$$L(x) = f(a) + f'(a)(x - a)$$

the linear approximation to f near a

The differential of
$$y = f(x)$$
 is

$$dy = f'(x) dx$$

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト

э

Functions of One Variable - Differentiability



Recall that if y = f(x), the *increment* of y as x changes from a to $a + \Delta x$ is

$$\Delta y = f(a + \Delta x) - f(a).$$

If f is differentiable at a, then

$$\Delta y = f'(a) \,\Delta x + \varepsilon \Delta x$$

where

$$\epsilon
ightarrow 0$$
 as $\Delta x
ightarrow 0$

That is, the linear approximation is very good as $\Delta x \rightarrow 0$.

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト

э

Derivatives - Two Variables

The derivatives $f_x(a, b)$ and $f_y(a, b)$ define a *tangent plane* to the graph of f at (a, b, f(a, b))

ヘロト ヘ週ト ヘヨト ヘヨト

æ



Derivatives - Two Variables

The derivatives $f_x(a, b)$ and $f_y(a, b)$ define a *tangent plane* to the graph of f at (a, b, f(a, b))

These derivatives define a linear function

$$L(x, y) = f(a, b)$$

+ $f_x(a, b)(x - a)$
+ $f_y(a, b)(x - b)$

the linear approximation to f near (a, b)

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト ・ ヨ



Derivatives - Two Variables

I

The derivatives $f_x(a, b)$ and $f_y(a, b)$ define a *tangent plane* to the graph of f at (a, b, f(a, b))

These derivatives define a linear function

$$L(x, y) = f(a, b)$$

+ $f_x(a, b)(x - a)$
+ $f_y(a, b)(x - b)$

the linear approximation to f near (a, b)

The differential of
$$z = f(x, y)$$
 is

$$dz = \frac{\partial f}{\partial x} \, dx + \frac{\partial f}{\partial y} \, dy$$

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト ・ ヨ



Find the Tangent Plane

If f has continuous partial derivatives, the tangent plane to z=f(x,y) at (a,b,f(a,b)) is

$$z - f(a, b) = f_x(a, b)(x - a) + f_y(a, b)(y - b)$$



1. Find the equation of the tangent plane to the surface

$$z = 2x^2 + y^2 - 5y$$

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト ・ ヨ

at (1, 2, -4).

Find the Tangent Plane

If f has continuous partial derivatives, the tangent plane to z=f(x,y) at (a,b,f(a,b)) is

$$z - f(a, b) = f_x(a, b)(x - a) + f_y(a, b)(y - b)$$



1. Find the equation of the tangent plane to the surface

$$z = 2x^2 + y^2 - 5y$$

at (1, 2, -4).

2. Find the equation of the tangent plane to the surface

$$z = e^{x-y}$$

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト ・ ヨ

at
$$(2, 2, 1)$$
.

The Tangent Plane Contains Tangent Lines



The red curves represent f(a, y) and f(x, b)

The blue lines are the tangent lines

$$r_1(t) = \langle a, b, f(a, b) \rangle + t \langle 1, 0, f_x(a, b) \rangle$$

$$r_2(t) = \langle a, b, f(a, b) \rangle + t \langle 0, 1, f_y(a, b) \rangle$$

Learning Goals

The Tangent Plane Defines a Linear Approximation



The tangent line is the graph of a linear function

$$L(x) = f(a) + f'(a)(x - a)$$

that approximates f(x) near x = a

The tangent plane is the graph of a linear function

L(x, y) = f(a, b) + $f_x(a, b)(x - a) + f_y(a, b)(y - b)$ that approximates f(x, y) near (x, y) =(a, b)

▲ロト ▲帰 ト ▲ ヨ ト ▲ ヨ ト ・ ヨ ・ の Q ()



The Linear Approximation

The linear approximation to f(x, y) at (a, b) is

$$L(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$$

- 1. Show that the linear approximation to $f(x, y) = e^x \cos(xy)$ at (0, 0) is L(x, y) = x + 1
- 2. Suppose that f(2,5) = 6, $f_x(2,5) = 1$, and $f_y(2,5) = -1$. Use a linear approximation to estimate f(2.2, 4.9)

Differentiability

If z = f(x, y), the *increment* of z as x changes from a to $a + \Delta x$ and y changes from b to $b + \Delta y$ is:

$$\Delta z = f(a + \Delta x, b + \Delta y) - f(a, b)$$

f is differentiable at (a, b) if

$$\Delta z = f_x(a, b)\Delta x + f_y(a, b)\Delta y + \varepsilon_1 \Delta x + \varepsilon_2 \Delta y$$

where ε_1 and ε_2 approach 0 as $(\Delta x, \Delta y) \rightarrow (0, 0)$.

Theorem If the partial derivatives f_x and f_y of f exist near (a, b), and are continuous at (a, b), then f is differentiable at (a, b).

1. Explain why the function $f(x, y) = \sqrt{xy}$ is differentiable at (1, 4) and find its linearization

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <

What Happens if f is not differentiable?



Use the definition to check that $f_x(0,0) = f_y(0,0) = 0$ Show that $f_x(x,y)$ and $f_y(x,y)$ are not continuous at (0,0)

◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへぐ

Differentials

For a function f of one variable, the differential of y = f(x) is given by

$$dy = f'(x) dx$$

For a function f of two variables, the differential of z = f(x, y) is

$$dz = f_x(x, y) \, dx + f_y(x, y) \, dy = \frac{\partial z}{\partial x} \, dx + \frac{\partial z}{\partial y} \, dy$$

- 1. The radius of a circle is measured as 10cm with an error of at most 0.2cm. What is the maximum calculated area of the circle?
- 2. The length and width of a rectangle are measured as 30cm and 24cm, with an error of at most 0.1cm each. What is the maximum error in the calculated area of the rectangle?

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 のへぐ

Three Variables

If w = f(x, y, z):

• The linear approximation of f at (a, b, c) is

$$L(x, y, z) = f(a, b, c) + f_x(a, b, c)(x - a) + f_y(a, b, c)(y - a) + f_z(a, b, c)(z - c)$$

• The *increment* of *w* is

$$\Delta w = f(x + \Delta x, y + \Delta y, z + \Delta z) - f(x, y, z)$$

• The differential dw is

$$dw = \frac{\partial w}{\partial x} \, dx + \frac{\partial w}{\partial y} \, dy + \frac{\partial w}{\partial z} \, dz$$

◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへぐ

Three Variables

The linear approximation of f at (a, b, c) is

$$L(x, y, z) = f(a, b, c) + f_x(a, b, c)(x - a) + f_y(a, b, c)(y - a) + f_z(a, b, c)(z - c)$$

Find the linear approximation to

$$f(x, y, z) = \sqrt{x^2 + y^2 + z^2}$$

at (x, y, z) = (3, 2, 6) and estimate

$$\sqrt{(3.02)^2 + (1.97)^2 + (5.99)^2}$$