▲□▶ ▲圖▶ ▲臣▶ ▲臣▶ ―臣 … のへで

Math 213 - Double Integrals Over Rectangles and Regions

Peter A. Perry

University of Kentucky

February 27, 2019

◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへで

Homework



- Webwork B7 on 14.8 (Lagrange multipliers) is due Friday
- Study for Quiz #6 on 14.6-14.7, tomorrow in recitation
- Re-read section 15.1, read section 15.2
- Begin work on problems 9-21 (odd), 27-43 (odd), 49 from 15.1

▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ

Unit II: Differential Calculus of Several Variables

- Lecture 12 Functions of Several Variables
- Lecture 13 Partial Derivatives
- Lecture 14 Tangent Planes and Linear Approximation
- Lecture 15 The Chain Rule
- Lecture 16 Directional Derivatives and the Gradient
- Lecture 17 Maximum and Minimum Values, I
- Lecture 18 Maximum and Minimum Values, II
- Lecture 19 Lagrange Multipliers
- Lecture 20 Double Integrals
- Lecture 21 Double Integrals over General Regions
- Lecture 22 Double Integrals in Polar Coordinates

Lecture 23 Exam II Review

Goals of the Day

- Understand why the volume under a surface can be computed as a double integral, a limit of double Riemann sums
- Understand how to compute double integrals over rectangles as *iterated integrals*
- Understand how to find the *average value* of a function of two variables over a rectangular domain

The area under the graph of y = f(x)between x = a and x = b is approximated by *Riemann sums*







▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ



A Riemann sum with n = 8 for $\int_{a}^{b} f(x) dx$

The area under the graph of y = f(x)between x = a and x = b is approximated by *Riemann sums*

$$\sum_{i=1}^n f(x_i^*) \, \Delta x$$

where

$$\Delta x = \frac{b-a}{n}$$
$$x_i = a + i\Delta x$$
$$x_i^* \in [x_{i-1}, x_i]$$

A Riemann sum with n = 8 for $\int_{a}^{b} f(x) dx$



▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ



The area under the graph of y = f(x)between x = a and x = b is approximated by *Riemann sums*

$$\sum_{i=1}^n f(x_i^*) \, \Delta x$$

where

$$\Delta x = \frac{b-a}{n}$$
$$x_i = a + i\Delta x$$
$$x_i^* \in [x_{i-1}, x_i]$$

A Riemann sum with n = 8 for $\int_{a}^{b} f(x) dx$

$$x_i^* = x_{i-1}$$
 (left endpoint)
 $x_i^* = x_i$ (right endpoint)

▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ



The area under the graph of y = f(x)between x = a and x = b is approximated by *Riemann sums*

$$\sum_{i=1}^n f(x_i^*) \, \Delta x$$

where

$$\Delta x = \frac{b-a}{n}$$
$$x_i = a + i\Delta x$$
$$x_i^* \in [x_{i-1}, x_i]$$

 $\begin{aligned} & x_i^* = x_{i-1} & \text{(left endpoint)} \\ & x_i^* = x_i & \text{(right endpoint)} \\ & x_i^* = \frac{x_{i-1} + x_i}{2} & \text{(midpoint)} \end{aligned}$



A Riemann sum with n = 8 for $\int_{a}^{b} f(x) dx$

▲□▶ ▲圖▶ ▲厘▶ ▲厘▶ 厘 の��

The exact area under the graph of f(x) between x = a and x = b is





The Fundamental Theorem of Calculus states that, if f is continuous on [a, b], then

$$\int_{a}^{b} f(x) \, dx = F(b) - F(a)$$

for any antiderivative F of f.

We extended these ideas to compute the *net* area under the graph of a signed function and the average value of a function f over an interval [a, b]

Calculus II: Volumes under Surfaces

S y х

Problem Find the volume between the rectangle

$$R = [a, b] \times [c, d]$$

in the xy plane and the surface

 $S = \{(x, y, z) : (x, y) \in R, z = f(x, y)\}$

if f is a continuous function.

- 1. Divide the rectangle R into an $n \times n$ 'grid' of subrectangles $R_{i,i}$
- 2. For each subrectangle $R_{i,j}$, make a box of height $f(x_i^*, y_i^*)$

3. Add up the volumes of the n^2 boxes

Learning Goals

Average Values

Volumes Under Surfaces





• Pick a rectangle $R_{i,j}$ in the grid, area ΔA



• Pick a rectangle $R_{i,j}$ in the grid, area ΔA

▲□▶ ▲圖▶ ▲臣▶ ▲臣▶ ―臣 … のへで

• Pick (x_i^*, y_i^*) in the rectangle



- Pick a rectangle $R_{i,j}$ in the grid, area ΔA
- Pick (x_i^*, y_i^*) in the rectangle
- Make a box of height f(x_i^{*}, y_j^{*}) over R_{i,j}

・ロト ・ 一下・ ・ モト ・ モト・

3



- Pick a rectangle $R_{i,j}$ in the grid, area ΔA
- Pick (x_i^*, y_i^*) in the rectangle
- Make a box of height f(x_i^{*}, y_j^{*}) over R_{i,j}

• The volume of the box is $V_{ij} = f(x_i^*, y_j^*) \Delta A$



- Pick a rectangle R_{i,j} in the grid, area ΔA
- Pick (x_i^*, y_i^*) in the rectangle
- Make a box of height $f(x_i^*, y_j^*)$ over $R_{i,j}$
- The volume of the box is $V_{ij} = f(x_i^*, y_j^*) \Delta A$
- The approximate volume under the surface is

 $\sum_{i=1}^{n}\sum_{j=1}^{n}f(x_{i}^{*},y_{j}^{*})\Delta A$

▲ロト ▲帰 ト ▲ ヨ ト ▲ ヨ ト ・ ヨ ・ の Q ()



- Pick a rectangle $R_{i,j}$ in the grid, area ΔA
- Pick (x_i^*, y_i^*) in the rectangle
- Make a box of height $f(x_i^*, y_j^*)$ over $R_{i,j}$
- The volume of the box is $V_{ij} = f(x_i^*, y_j^*) \Delta A$
- The approximate volume under the surface is

$$\sum_{i=1}^{n}\sum_{j=1}^{n}f(x_{i}^{*},y_{j}^{*})\Delta A$$

▲ロト ▲帰 ト ▲ ヨ ト ▲ ヨ ト ・ ヨ ・ の Q ()

$$V = \lim_{n \to \infty} \sum_{i,j=1}^{n} f(x_i^*, y_j^*) \Delta A$$

Volumes By Integration

If $R = [a, b] \times [c, d]$ and

$$S = \{(x, y, z) : z = f(x, y), (x, y) \in R\}$$

then the volume between R and S is

$$V = \lim_{n \to \infty} \sum_{i,j=1}^{n} f(x_i^*, y_j^*) \Delta A = \iint_R f(x, y) \, dA$$



Find
$$\iint_R f(x, y) \, dA$$
 if

$$R = [-1,1] \times [0,2]$$

and

$$f(x,y) = \sqrt{1-x^2}$$

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <

Can you do this without calculus?

Iterated Integrals

When you studied one-variable calculus, you first found out how to compute antiderivatives and then you learned how to compute definite integrals using them

Now that you're studying two variable calculus, you'll first learn about *iterated integrals* and then learn how to compute integrals over rectangles with them.

Suppose *R* is a rectangle $[a, b] \times [c, d]$ and *f* is a continuous function on *R*. Then

$$A(x) = \int_{c}^{d} f(x, y) \, dy$$

is a function of x. For example if $f(x, y) = x^2 y$ and $R = [1, 2] \times [3, 4]$, then

$$\int_{3}^{4} (x^{2}y) \, dy = \left. \frac{x^{2}y^{2}}{2} \right|_{3}^{4} = \frac{7}{2}x^{2}$$

We then compute $\int_{a}^{b} A(x) dx$. For example

$$\int_{1}^{2} \frac{7}{2} x^{2} dx = \left. \frac{7}{6} x^{3} \right|_{x=1}^{x=2} = \frac{49}{6}$$

<ロト 4 回 ト 4 回 ト 4 回 ト 1 回 9 Q Q</p>

Iterated Integrals

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <

Average Values

Iterated Integrals

If f(x, y) is a continuous function on a rectangle $R = [a, b] \times [c, d]$, the **iterated integral** of f is

$$\int_{a}^{b} \int_{c}^{d} f(x, y) \, dy \, dx = \int_{a}^{b} \left(\int_{c}^{d} f(x, y) \, dy \right) \, dx$$

1. Find
$$\int_{1}^{4} \int_{0}^{2} (6x^{2}y - 2x) dy dx$$

2. Find $\int_{1}^{3} \int_{1}^{5} \frac{\ln y}{xy} dy dx$
3. (Ringer) $\int_{0}^{1} \int_{1}^{2} (x + e^{-y}) dx dy$

Fubini's Theorem

Theorem If *f* is continuous on the rectangle

$$R = \{(x, y) : a \le x \le b, \ c \le y \le d\}$$

then

$$\iint_R f(x,y) \, dA = \int_a^b \int_c^d f(x,y) \, dy \, dx = \int_c^d \int_a^b f(x,y) \, dx \, dy.$$

Evaluate these double integrals.

1.
$$\iint_{R} x \sec^{2} y \, dA, \ R = [0, 2] \times [0, \pi/4]$$

2.
$$\iint_{R} \frac{xy^{2}}{x^{2}+1} \, dA, \ R = [0, 1] \times [-3, 3]$$

3.
$$\iint_{R} \frac{1}{1+x+y} \, dA, \ R = [1, 3] \times [1, 2]$$

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <

Volumes by Iterated Integrals

1. Find the volume of the solid that lies under the plane

4x + 6y - 12z + 15 = 0

and above the rectangle $[-1,2]\times [-1,1]$

2. Find the volume of the solid lying under the elliptic paraboloid

 $x^2/4 + y^2/9 + z = 1$

and above the rectangle $[-1,1]\times [-2,2]$

Preview: Integrals over General Regions

Find

if

y y = 4x/3y = x/31 3 x

$$\iint_D (x+3y) \, dA$$

$$D = \{1 \le x \le 3, x/3 \le y \le 4x/3\}$$

◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへぐ

Preview: Integrals over General Regions

if



Find

$$\iint_{D} (x + 3y) \, dA$$
if

$$D = \{1 \le x \le 3, \ x/3 \le y \le 4x/3$$

$$\iint_{D} (x+3y) \, dA =$$
$$\int_{1}^{3} \left(\int_{x/3}^{4x/3} (x+3y) \, dy \right) \, dx$$

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 のへぐ

}

Learning Goals From Areas to Volumes Iterated Integrals General Regions Average Values







if D is the region enclosed by the curves

$$x = 2 - y^2$$

and

$$x = -2 + y^2.$$

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 のへぐ

Learning Goals	From Areas to Volumes	Iterated Integrals	General Regions	Average Values



$$\iint_D 1 \, dA$$

if D is the region enclosed by the curves

$$x = 2 - y^2$$

and

Find

 $x = -2 + y^2.$

$$D = \{-\sqrt{2} \le y \le \sqrt{2}, \ -2 + y^2 \le x \le 2 + y^2\}$$

Learning Goals From Areas to Volumes Iterated Integrals General Regions Average Values



$$\iint_D 1 \, dA$$

if D is the region enclosed by the curves

$$x = 2 - y^2$$

and

Find

 $x = -2 + y^2.$

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 のへぐ

$$D = \{-\sqrt{2} \le y \le \sqrt{2}, -2 + y^2 \le x \le 2 + y^2\}$$
$$\iint_D 1 \, dA = \int_{-\sqrt{2}}^{\sqrt{2}} \left(\int_{-2 + y^2}^{2 - y^2} 1 \, dx\right) \, dy$$

Average Values

The average value of y = f(x) on [a, b] is

f

$$f_{\rm av} = \frac{1}{b-a} \int_a^b f(x) \, dx$$

The average value of z = f(x, y) on a rectangle R with area A(R) is

$$f_{\rm av} = \frac{1}{A(R)} \iint_R f(x, y) \, dA$$



Find the average value of $f(x, y) = x^2 y$ over a rectangle with vertices (-1, 0), (-1, 5), (1, 5), and (1, 0).