

# Math 213 - Change of Variables, Part I

Peter A. Perry

University of Kentucky

March 25, 2019

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 の�?



- Re-read section 15.9 for Wednesday
- Practice problems for 15.9 are 1-19 (odd), 21, 23
- Webwork C4 on 15.8 is due on Wednesday Night

▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ

• You have a quiz on 15.7-15.8 on Thursday

▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ

# Unit III: Integral Calculus, Vector Fields

- Lecture 24 Triple Integrals
  Lecture 25 Triple Integrals, Continued
  Lecture 26 Triple Integrals Cylindrical Coordinates
  Lecture 27 Triple Integrals Spherical Coordinates
  Lecture 28 Change of Variables for Multiple Integrals, I
  Lecture 29 Change of Variable for Multiple Integrals, II
- Lecture 30 Vector Fields
- Lecture 31 Line Integrals (Scalar Functions)
- Lecture 32 Line Integrals (Vector Functions)
- Lecture 33 Fundamental Theorem for Line Integrals
- Lecture 34 Green's Theorem

Lecture 35 Exam III Review



- Understand what a transformation  $\mathcal{T}$  between two regions in the plane is
- Understand how to compute the *Jacobian Matrix* and *Jacobian determinant* of a transformation and understand what the Jacobian determinant measures
- Understand how to compute double integrals using the change of variables formula

Change of Variables Formula

# Preview: Calculus I versus Calculus III

If x = g(u) maps [c, d] to [a, b], then

$$\int_a^b f(x) \, dx = \int_c^d f(g(u)) \, g'(u) \, du$$

In other words,

$$\int_{a}^{b} f(x) \, dx = \int_{c}^{d} f(x(u)) \, \frac{dx}{du} \, du$$

If x = g(u, v), y = h(u, v), and if the region S in the uv plane is mapped to the region R in the xy plane, then

$$\iint_{R} f(x, y) \, dA = \iint_{S} f(x(u, v), y(u, v)) \, J(u, v) \, du \, dv$$

The Jacobian determinant

$$J(u,v) = \left|\frac{\partial(x,y)}{\partial(u,v)}\right|$$

measures how areas change under the map  $(u, v) \mapsto (x, y)$ .

Preview

Transformations

The Jacobiar

Change of Variables Formula

# Transformations



The polar coordinate map

 $x = r \cos \theta$ ,  $y = r \sin \theta$ 

defines a transformation  $T: S \rightarrow R$ 

Before, we called R a *polar rectangle* 

Here are the corresponding sides of the rectangle in the  $r\theta$  plane and the polar rectangle in the *xy* plane:

イロト 不得 トイヨト イヨト

3



Preview

Transformations

The Jacobiar

Change of Variables Formula

# Transformations





 $x = r \cos \theta$ ,  $y = r \sin \theta$ 

defines a transformation  $T: S \rightarrow R$ 

Before, we called R a *polar rectangle* 

Here are the corresponding sides of the rectangle in the  $r\theta$  plane and the polar rectangle in the *xy* plane:



• The line r = a

Preview

Transformations

The Jacobiar

Change of Variables Formula

# Transformations



R

 $\theta = \alpha$ 

х

r = a

The polar coordinate map

 $x = r \cos \theta$ ,  $y = r \sin \theta$ 

defines a transformation  $T: S \rightarrow R$ 

Before, we called R a *polar rectangle* 

Here are the corresponding sides of the rectangle in the  $r\theta$  plane and the polar rectangle in the *xy* plane:



• The line  $\theta = \alpha$ 

Preview

Transformations

The Jacobiar

Change of Variables Formula

# Transformations



The polar coordinate map

 $x = r \cos \theta$ ,  $y = r \sin \theta$ 

defines a transformation  $T: S \rightarrow R$ 

Before, we called R a *polar rectangle* 

Here are the corresponding sides of the rectangle in the  $r\theta$  plane and the polar rectangle in the *xy* plane:

- The line r = a
- The line  $\theta = \alpha$
- The line r = b

Preview

Transformations

The Jacobiar

Change of Variables Formula

# Transformations



The polar coordinate map

 $x = r \cos \theta$ ,  $y = r \sin \theta$ 

defines a transformation  $T: S \rightarrow R$ 

Before, we called R a *polar rectangle* 

Here are the corresponding sides of the rectangle in the  $r\theta$  plane and the polar rectangle in the *xy* plane:

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト ・ ヨ

- The line r = a
- The line  $\theta = \alpha$
- The line r = b
- The line  $\theta = \beta$

Preview

Transformations

The Jacobiar

Change of Variables Formula

# Transformations



The polar coordinate map

 $x = u \cos v$ ,  $y = u \sin v$ 

defines a transformation  $T: S \rightarrow R$ 

Before, we called R a *polar rectangle* 

Here are the corresponding sides of the rectangle in the  $u\theta$  plane and the polar rectangle in the *xy* plane:



Preview

Transformations

The Jacobiar

Change of Variables Formula

# Transformations





 $x = u \cos v$ ,  $y = u \sin v$ 

defines a transformation  $T: S \rightarrow R$ 

Before, we called R a *polar rectangle* 

Here are the corresponding sides of the rectangle in the  $u\theta$  plane and the polar rectangle in the *xy* plane:

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト ・ ヨ



• The line u = a

Preview

Transformations

The Jacobiar

Change of Variables Formula

# Transformations





The polar coordinate map

 $x = u \cos v$ ,  $y = u \sin v$ 

defines a transformation  $T: S \rightarrow R$ 

Before, we called R a *polar rectangle* 

Here are the corresponding sides of the rectangle in the  $u\theta$  plane and the polar rectangle in the *xy* plane:

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト ・ ヨ

- The line u = a
- The line  $v = \alpha$

Preview

Transformations

The Jacobiar

Change of Variables Formula

# Transformations



The polar coordinate map

 $x = u \cos v$ ,  $y = u \sin v$ 

defines a transformation  $T: S \rightarrow R$ 

Before, we called R a *polar rectangle* 

Here are the corresponding sides of the rectangle in the  $u\theta$  plane and the polar rectangle in the *xy* plane:

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト ・ ヨ

- The line u = a
- The line  $v = \alpha$
- The line u = b

Preview

Transformations

The Jacobiar

Change of Variables Formula

# Transformations



The polar coordinate map

 $x = u \cos v$ ,  $y = u \sin v$ 

defines a transformation  $T: S \rightarrow R$ 

Before, we called R a *polar rectangle* 

Here are the corresponding sides of the rectangle in the  $u\theta$  plane and the polar rectangle in the *xy* plane:

◆□▶ ◆□▶ ◆□▶ ◆□▶ ●□

- The line u = a
- The line  $v = \alpha$
- The line u = b
- The line  $v = \beta$

Preview

Transformations

The Jacobiar

Change of Variables Formula

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 の�?

#### Transformations

The equations

$$x = u^2 - v^2, \quad y = 2uv$$

defines a transformation  $T: S \rightarrow R$ 





Preview

Transformations

The Jacobiar

Change of Variables Formula

◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへぐ

### Transformations

The equations

$$x = u^2 - v^2, \quad y = 2uv$$

defines a transformation  $T: S \rightarrow R$ 

•  $v = 0, 0 \le u \le 1$  maps to  $0 \le x \le 1, y = 0$ 





Preview

Transformations

The Jacobiar

Change of Variables Formula

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <

#### Transformations

The equations

$$x = u^2 - v^2, \quad y = 2uv$$

defines a transformation  $T: S \rightarrow R$ 

- $v = 0, 0 \le u \le 1$  maps to  $0 \le x \le 1, y = 0$
- u = 1,  $0 \le v \le 1$  maps to the parametric curve  $x = 1 v^2$ , y = 2v





Preview

Transformations

The Jacobiar

Change of Variables Formula

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <

#### Transformations

The equations

$$x = u^2 - v^2, \quad y = 2uv$$

defines a transformation  $T: S \rightarrow R$ 

- $v = 0, 0 \le u \le 1$  maps to  $0 \le x \le 1, y = 0$
- u = 1,  $0 \le v \le 1$  maps to the parametric curve  $x = 1 v^2$ , y = 2v
- $v = 1, 0 \le u \le 1$  maps to the parametric curve  $x = u^2 - 1, y = 2u$





Preview

Transformations

The Jacobian

Change of Variables Formula

#### Transformations

The equations

$$x = u^2 - v^2, \quad y = 2uv$$

defines a transformation  $T: S \rightarrow R$ 

- $v = 0, 0 \le u \le 1$  maps to  $0 \le x \le 1, y = 0$
- u = 1,  $0 \le v \le 1$  maps to the parametric curve  $x = 1 v^2$ , y = 2v
- v = 1,  $0 \le u \le 1$  maps to the parametric curve  $x = u^2 1$ , y = 2u

• 
$$u = 0, 0 \le v \le 1$$
 maps to  
 $-1 \le x \le 0, y = 0$ 





◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへぐ



1. Find the image of

$$S = \{(u, v) : 0 \le u \le 3, \ 0 \le v \le 2\}$$

under the transformation x = 2u + 3v, y = u - v

2. Find the image of the disc  $u^2 + v^2 \le 1$  under the transformation x = au, y = bv

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 の�?

# The Jacobian

In one variable calculus, the way a transformation x = g(u) changes lengths of interals is measured by g'(u):

 $\Delta x = g'(u)\Delta u$ 



In two variable calculus, the way a transformation

$$x = g(u, v), \quad y = h(u, v)$$

changes areas is measured by the Jacobian determinant

$$J(u, v) = \begin{vmatrix} \partial x / \partial u & \partial x / \partial v \\ 0 & 0 \\ \partial y / \partial u & \partial y / \partial v \end{vmatrix}, \quad \Delta A = |J(u, v)| \Delta u \Delta v$$

We'll now see why this is the case...

◆□ > ◆□ > ◆豆 > ◆豆 > ̄豆 \_ のへぐ

ヘロン 人間 とくほと くほとう

æ

# The Jacobian

A transformation x = g(u, v), y = h(u, v) maps a small rectangle S into a distorted rectangle R through the rule

$$\mathbf{r}(u, v) = g(u, v)\mathbf{i} + h(u, v)\mathbf{j}$$



#### The Jacobian

A transformation x = g(u, v), y = h(u, v) maps a small rectangle S into a distorted rectangle R through the rule

$$\mathbf{r}(u, v) = \mathbf{g}(u, v)\mathbf{i} + h(u, v)\mathbf{j}$$



*R* has approximate area  $|\mathbf{r}_u \times \mathbf{r}_v| \Delta u \Delta v$  where

$$\mathbf{r}_{u} = \frac{\partial x}{\partial u}\mathbf{i} + \frac{\partial y}{\partial u}\mathbf{j}, \qquad \mathbf{r}_{v} = \frac{\partial x}{\partial v}\mathbf{i} + \frac{\partial y}{\partial v}\mathbf{j}$$

◆□ > ◆□ > ◆豆 > ◆豆 > ̄豆 = のへで



*R* has approximate area  $|\mathbf{r}_u \times \mathbf{r}_v| \Delta u \Delta v$  where

$$\mathbf{r}_{u} = \frac{\partial x}{\partial u}\mathbf{i} + \frac{\partial y}{\partial u}\mathbf{j}, \qquad \mathbf{r}_{v} = \frac{\partial x}{\partial v}\mathbf{i} + \frac{\partial y}{\partial v}\mathbf{j}$$

Compute

$$\mathbf{r}_{u} \times \mathbf{r}_{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial x}{\partial u} & \frac{\partial y}{\partial u} & \mathbf{0} \\ \frac{\partial x}{\partial v} & \frac{\partial y}{\partial v} & \mathbf{0} \end{vmatrix} = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$

・ロト ・四ト ・ヨト ・ヨト

æ



x

・ロト ・ 一 ト ・ モト ・ モト

æ

$$dA \simeq \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \Delta u \, \Delta v$$

where

$$\frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \partial x/\partial u & & \partial x/\partial v \\ \partial y/\partial u & & \partial y/\partial v \end{vmatrix}$$

is the Jacobian determinant of the transformation

The area of R is approximately

и

Learning Goals	Preview	Transformations	The Jacobian	Change of Variables Formula
The Jacobian				

◆□ ▶ < 圖 ▶ < 圖 ▶ < 圖 ▶ < 圖 • 의 Q @</p>

Find the Jacobian determinant of the following transformations.

1. 
$$x = 2u + 3v$$
,  $y = u - v$   
2.  $x = au$ ,  $y = bv$   
3.  $x = u^2 - v^2$ ,  $y = 2uv$ 

Change of Variables Formula

### Area Change in Polar Coordinates

Consider the transformation  $x = u \cos v$ ,  $y = u \sin v$ 



so



The Jacobian Determinant of a transformation x = g(u, v), y = h(u, v) is denoted

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}$$

The notation

$$\left|\frac{\partial(x,y)}{\partial(u,v)}\right|$$

◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへぐ

denotes the absolute value of this determinant.

# Change of Variables Formula

If the transformation x = g(u, v), y = h(u, v) maps a region S in the *uv*-plane to a region R in the xy plane:

$$\iint_{R} f(x, y) dA \simeq \sum_{i=1}^{n} \sum_{j=1}^{n} f(x_{i}, y_{j}) \Delta A$$
$$\simeq \sum_{i=1}^{n} \sum_{j=1}^{n} f(g(u_{i}, v_{j}), h(u_{i}, v_{j})) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \Delta u \Delta v$$
$$\iint_{S} f(g(u, v), h(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

**Change of Variables in a Double Integral** If *T* is a one-to-one transformation with nonzero Jacobian and  $T: S \rightarrow R$ , then

$$\iint_{R} f(x, y) \, dA = \iint_{S} f(x(u, v), y(u, v)) \, \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du \, dv$$

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ ─臣 ─のへで

# Change of Variables Formula

**Change of Variables in a Double Integral** If *T* is a one-to-one transformation with nonzero Jacobian and  $T: S \rightarrow R$ , then

$$\iint_{R} f(x, y) \, dA = \iint_{S} f(x(u, v), y(u, v)) \, \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du \, dv$$

- 1. Use the transformation x = 2u + v, y = u + 2v to find  $\iint_R (x 3y) dA$  if *R* is the triangular region with vertices (0, 0), (2, 1) and (1, 2)
- 2. Find  $\iint_R (x+y)e^{x^2-y^2} dA$  if R is the rectangle enclosed by x-y=0, x-y=2, x+y=0, and x+y=3.