◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 のへぐ

Math 213 - Line Integrals I

Peter A. Perry

University of Kentucky

April 1, 2019



▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ

Homework



• Work on Stewart problems for 16.2: 1-21 (odd), 33-41 (odd), 49, 50

• Webwork C1 on section 16.1 is due Wednesday

▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ

Unit IV: Vector Calculus

- Lecture 24Triple IntegralsLecture 25Triple Integrals, ContinuedLecture 26Triple Integrals Cylindrical CoordinatesLecture 27Triple Integrals Spherical CoordinatesLecture 28Change of Variables for Multiple Integrals, ILecture 29Change of Variable for Multiple Integrals, II
- Lecture 30 Vector Fields
- Lecture 31 Line Integrals (Scalar Functions)
- Lecture 32 Line Integrals (Vector Functions)
- Lecture 33 Fundamental Theorem for Line Integrals
- Lecture 34 Green's Theorem

Lecture 35 Exam III Review

Learning Goals

Overview

◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへぐ

Goals of the Day

- Know how to compute line integrals of a scalar function in the plane
- Know how to compute line integrals of a scalar function in space

Preview: Line Integrals

Our next topic will be integrals of *scalar functions* and *vector functions* over curves in the plane and in space. If C is a curve in the plane or in space, we'll learn how to compute:

- $\int_C f(x, y) ds$, the integral of a scalar function over a plane curve C
- $\int_C f(x, y, z) \, ds$, the integral of a scalar function over a space curve C
- $\int_C \mathbf{F} \cdot d\mathbf{r}$, the integal of a vector function $\mathbf{F}(x, y)$ over a plane curve C
- $\int_C \mathbf{F} \cdot d\mathbf{r}$, the integral of a vector function $\mathbf{F}(x, y, z)$ over a space curve C

In all cases, we'll reduce these to Calculus I and II type integrals by parameterizing the curve C. We'll also learn how to compute integrals like

- $\int_C f(x, y) dx$
- $\int_C f(x, y) dy$

Parameterizing Paths



Parameterize the following paths:

- 1. The first planar path shown on the left
- 2. The second planar path shown on the left
- 3. The path connecting $(\mathbf{0},\mathbf{0},\mathbf{0})$ to $(\mathbf{1},\mathbf{0},\mathbf{1})$
- 4. The path connecting (1, 0, 1) to (1, 2, 0)

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト ・ ヨ

The Integral of a Scalar Function over a Plane Curve

If C is a plane curve, the **line integral of** f **along** C is

$$\int_C f(x, y) \, ds = \lim_{n \to \infty} \sum_{i=1}^n f(x_i^*, y_i^*) \, \Delta s_i$$

where we approximate the curve by n line segments of length Δs_i

As a practical matter, if C is parameterized by (x(t), y(t)) for $a \le t \le b$,

$$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

so

$$\int_{C} f(x, y) \, ds = \int_{a}^{b} f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^{2} + \left(\frac{dy}{dt}\right)^{2}} \, dt$$

▲□▶ ▲圖▶ ▲≣▶ ▲≣▶ = 三 - のへ⊙

The Integral of a Scalar Function over a Plane Curve

if C is parameterized by (x(t), y(t)) for $a \leq t \leq b$, then

$$\int_{C} f(x, y) \, ds = \int_{a}^{b} f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^{2} + \left(\frac{dy}{dt}\right)^{2}} \, dt$$

1. Find $\int_C (x/y) ds$ if C is the curve $x = t^2$, y = 2t for $0 \le t \le 3$ 2. Find $\int_C xy^4 ds$ if C is the right half of the circle $x^2 + y^2 = 16$

Line Integrals over Piecewise Smooth Curves



A curve *C* is *piecewise smooth* if it is a union of smooth curves C_1, \ldots, C_n . Some examples are shown at left.

If ${\it C}$ consists of seveal smooth components, then

$$\int_C f(x, y) \, ds = \sum_{i=1}^n \int_{C_i} f(x, y) \, ds$$

Notice that each of these curves has an *orientation* that determines how the curve is parameterized—the parameterization should "follow the arrows."

1. Find $\int_C xy \, ds$ if C is the first curve shown at left.

▲ロト ▲帰 ト ▲ ヨ ト ▲ ヨ ト ・ ヨ ・ の Q ()

Another Kind of Line Integral

For later use, we'll also need the line integral of f with respect to x and the line integral of f with respect to y:

$$\int_{C} f(x, y) dx = \int_{a}^{b} f(x(t), y(t)) x'(t) dt$$
$$\int_{C} f(x, y) dy = \int_{a}^{b} f(x(t), y(t)) y'(t) dt$$

- 1. Find $\int_C e^x dx$ if C is the arc of the curve $x = y^3$ from (-1, -1) to (1, 1)
- 2. Find $\int_C x^2 dx + y^2 dy$ if C is the arc of the circle $x^2 + y^2 = 4$ from (2,0) to (0,2) followed by the line segment from (0,2) to (4,3)

・ロト ・ 西ト ・ モト ・ モー ・ つへぐ

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <

Summary of Line Integrals in the Plane

If C is a parameterized curve (x(t), y(t)) where $a \le t \le b$:

$$\int_C f(x, y) dx = \int_a^b f(x(t), y(t)) x'(t) dt$$
$$\int_C f(x, y) dy = \int_a^b f(x(t), y(t)) y'(t) dt$$
$$\int_C f(x, y) ds = \int_a^b f(x(t), y(t)) \sqrt{(x'(t)^2 + y'(t)^2} dt$$

Applications - Center of Mass

A wire of mass *m* and density $\rho(x, y)$ along a curve *C* has center of mass

$$\overline{x} = \frac{1}{m} \int_C x \rho(x, y) \, ds$$
$$\overline{y} = \frac{1}{m} \int_C y \rho(x, y) \, ds$$



A thin wire has the shape of the first quadrant part of a circle with center at the origin and radius *a*. If the density of the wire is

$$\rho(x,y) = kxy,$$

find the mass and center of mass of the wire.

Line Integrals in Space

If C is a space curve (x(t), y(t), z(t)) where $a \le t \le b$, then

$$\int_{C} f(x, y, z) ds = \int_{a}^{b} f(x(t), y(t), z(t)) \sqrt{(x'(t))^{2} + (y'(t))^{2} + (z'(t))^{2}} dt$$

1. Find $\int_C (x^2 + y^2 + z^2) ds$ if C is the space curve $(x(t), y(t), z(t)) = (t, \cos 2t, \sin 2t)$ for $0 \le t \le 2\pi$

More Line Integrals in Space

Can you guess how to define $\int_C f(x, y, z) dx$, $\int_C f(x, y, z) dy$, and $\int_C f(x, y, z) dz$?

1. Find $\int_C (x+z) dx + \int_C (x+z) dy + \int_C (x+y) dz$ if C consists of the line segments from (0,0,0) to (1,0,1) and from (1,0,1) to (0,1,2)

