◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 の�?

# Math 213 - Divergence and Curl

#### Peter A. Perry

University of Kentucky

April 12, 2019



The Big Three

▲□▶ ▲圖▶ ▲臣▶ ▲臣▶ = 臣 = のへで





 Work on Stewart problems for 16.5: 1-11 (odd), 12, 13-17 (odd), 21, 23, 25 Curl

Divergence

The Big Three

▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ

#### Unit IV: Vector Calculus

- Lecture 36 Curl and Divergence Lecture 37 Parametric Surfaces Lecture 38 Surface Integrals Lecture 39 Stokes' Theorem Lecture 40 The Divergence Theorem
- Lecture 41 Final Review, Part I Lecture 42 Final Review, Part II

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <

# Goals of the Day

This lecture is about two very important 'derivatives' of a vector field. You'll learn:

- How to compute the *curl* of a vector field and what it measures
- How to compute the *divergence* of a vector field and what it measures
- (Sneak preview) The theorems that give the meaning of divergence and curl

# Curl

If  $F = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$  is a vector field on  $\mathbb{R}^3$ , and the partial derivatives of P, Q, and R all exist, then the *curl* of  $\mathbf{F}$  is a new vector field:

$$\operatorname{curl} \mathbf{F} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z}\right) \mathbf{i} + \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x}\right) \mathbf{j} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) \mathbf{k}$$

This new vector field measures the "rotation" of the vector field **F** at a given point (x, y, z):

- Its direction is the axis of rotation, dictated by the right-hand rule
- Its *magnitude* is the angular speed of rotation



$$\operatorname{curl} \mathbf{F} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z}\right) \mathbf{i} + \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x}\right) \mathbf{j} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) \mathbf{k}$$

The Big Three

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 の�?

The new vector field curl  ${\bf F}$  is sometimes written  $\nabla\times {\bf F}$  because of an easier-to-remember formula:

$$\operatorname{curl} \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix}$$

Learning Goals

Curl

Divergence

The Big Three

#### Curl



If  $\mathbf{F} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$  then  $\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x & y & z \end{vmatrix} = \mathbf{0}$ 

 $\mathbf{F}(x, y, z) = y\mathbf{i} - x\mathbf{j} + 0\mathbf{k}$ 



If  $\mathbf{F} = y\mathbf{i} - x\mathbf{j} + 0\mathbf{k}$  then  $\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial x} & \frac{\partial}{\partial x} \end{vmatrix}$ 

$$= \begin{vmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y & -x & 0 \end{vmatrix} = -2\mathbf{k}$$

▲□▶ ▲圖▶ ▲臣▶ ▲臣▶ ―臣 … のへで



A gradient vector field has zero curl:

$$\nabla \times (\nabla f) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \end{vmatrix}$$
$$= \left(\frac{\partial^2 f}{\partial x \partial y} - \frac{\partial^2 f}{\partial y \partial x}\right) \mathbf{i} + \left(\frac{\partial^2 f}{\partial y \partial z} - \frac{\partial^2 f}{\partial z \partial y}\right) \mathbf{j} + \left(\frac{\partial^2 f}{\partial z \partial x} - \frac{\partial^2 f}{\partial x \partial z}\right) \mathbf{k}$$
$$= \mathbf{0}$$

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 の�?

so the curl "detects" conservative vector fields.

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <

## Divergence

The *divergence* of a vector field  $\mathbf{F} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$  is a *scalar* function:

div 
$$\mathbf{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

Sometimes div **F** is written  $\nabla \cdot \mathbf{F}$ :

$$\nabla \cdot \mathbf{F} = \left\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\rangle \cdot \left\langle P, Q, R \right\rangle = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

The divergence computes the outflow per unit volume of the vector field (thought of as a velocity field)

## Divergence



If  $\mathbf{F} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$  then  $\nabla \cdot \mathbf{F} = \frac{\partial x}{\partial x} + \frac{\partial y}{\partial y} + \frac{\partial z}{\partial z} = 3$ 

(same outflow at each point of space)

 $\mathbf{F}(x, y, z) = y\mathbf{i} - x\mathbf{j} + 0\mathbf{k}$ 



If 
$$\mathbf{F} = y\mathbf{i} - x\mathbf{j} + 0\mathbf{k}$$
 then  
 $\nabla \cdot \mathbf{F} = \frac{\partial y}{\partial x} - \frac{\partial x}{\partial y} + 0 = 0$ 

(no outflow anywhere in space)

◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへぐ

◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへぐ

## Divergence

Remember that  $\nabla\times(\nabla f)=$  0? There is an analogous result for the divergence:

 $\operatorname{div}\operatorname{curl} \boldsymbol{F}=0$ 

You can see this using the definitions of divergence and curl:

div curl 
$$\mathbf{F} = \frac{\partial}{\partial x} \left( \frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) + \frac{\partial}{\partial y} \left( \frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} \right) + \frac{\partial}{\partial z} \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right)$$

The second partial derivatives cancel in pairs by Clairaut's theorem.

It turns out that any vector field F can be written as

$$\mathbf{F} = \nabla f + \nabla \times \mathbf{A}$$

for a scalar potential f and a vector potential A.

▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ

## Divergence and Curl

If f is a scalar function and **F** is a vector function, which of these expressions make sense? Do they define a scalar or a vector? Remember that

- curl **F** is a vector
- div F is a scalar
- (a)  $\operatorname{curl} f$  (b)  $\operatorname{grad} f$
- (c) div **F** (d) curl(grad f)
- $(e) \quad \mathsf{grad}\, \mathbf{F} \qquad \qquad (f) \quad \mathsf{grad}(\mathsf{div}\, \mathbf{F}) \\$
- (g)  $\operatorname{div}(\operatorname{grad} f)$  (h)  $\operatorname{grad}(\operatorname{div} f)$
- (i)  $\operatorname{curl}(\operatorname{curl} \mathbf{F})$  (j)  $\operatorname{div}(\operatorname{div} f)$

Learning Goals

Curl

Divergence

The Big Three

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 の�?

#### Conservative Vector Fields Again

Determine whether the vector field

$$F(x, y, z) = y^2 z^3 i + 2xyz^3 j + 3xy^2 z^2 k$$

is conservative and, if so, find a function f so that  $\nabla f = \mathbf{F}$ .

Learning Goals

Cur

Divergence

The Big Three

#### Vector Identities

Show that  $\operatorname{div}(f\mathbf{F}) = \nabla f \cdot \mathbf{F} + f \operatorname{div} \mathbf{F}$ 



# Divergence Theorem, Stokes' Theorem

**Divergence Theorem** Suppose E is a simple solid region and S is its boundary. Let **N** be the outward normal to S. Then

$$\iint_{S} \mathbf{F} \cdot \mathbf{N} \, dS = \iiint_{E} \operatorname{div} \mathbf{F} \, dV$$

**Stokes' Theorem** Suppose *S* is an oriented piecewise-smooth surface with outward normal N, bounded by a simple closed curve *C* with piecewise smooth boundary. Then

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \iint_S \operatorname{curl} \mathbf{F} \cdot \mathbf{N} \, dS$$

# The 'Big Three' Theorems of Vector Calculus

$$F(b) - F(a) = \int_{a}^{b} F'(x) \, dx \qquad (\text{Fundamental})$$

$$\int_{C} P \, dx + Q \, dy = \iint_{D} \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \, dA \tag{Green}$$

$$\int_{C} \mathbf{F} \cdot d\mathbf{r} = \iint_{S} \operatorname{curl} \mathbf{F} \cdot \mathbf{N} \, dS \tag{Stokes}$$

$$\iint_{S} \mathbf{F} \cdot \mathbf{N} \, dS = \iiint_{E} \operatorname{div} \mathbf{F} \, dV \qquad \text{(Divergence)}$$