Review

Scalar Surface Integrals

Oriented Surfaces

Vector Surface Integrals

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 のへぐ

Math 213 - Surface Integrals

Peter A. Perry

University of Kentucky

April 17, 2019

Vector Surface Integrals

Final Exam

Your **Final Exam** is on **Wednesday**, **May 1 at 6:00 PM**. Please be at least five minutes early and bring your student ID. Ground rules are the same as for previous exams. Make a note of your exam room:

Section	Instructor	Exam Room
Sections 001, 002	(Kasey Bray)	CB 102
Section 003	(Deborah Wilkerson)	CB 110

Vector Surface Integrals

・ロト ・ 理 ト ・ ヨ ト ・ ヨ ト ・ ヨ

Homework

- Homework D2 is due tonight
- Work on Stewart problems for 16.7: 5-19 alternate odd, 21-31 alternate odd
- Study for your last quiz of the semester, Quiz 10, on sections 16.5-16.6 (divergence and curl, parameterized surfaces)
- Read section 16.8 for Wednesday, Aprll 17

Vector Surface Integrals

▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ

Unit IV: Vector Calculus

- Lecture 36 Curl and Divergence Lecture 37 Parametric Surfaces Lecture 38 Surface Integrals Lecture 39 Stokes' Theorem Lecture 40 The Divergence Theorem Lecture 41 Final Review, Part I
- Lecture 42 Final Review, Part II

Vector Surface Integrals

< □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > <

Goals of the Day

This lecture is about parametric surfaces. You'll learn:

- How to integrate a scalar function over a parameterized surface
- What an *oriented surface* is and how to compute its *unit normal*
- How to integrate a vector field over a parameterized surface

Sneak Preview - Scalar Surface Integrals

Scalar Line Integrals

If C is parameterized by

$$\mathbf{r}(t) = \langle \mathbf{x}(t), \mathbf{y}(t), \mathbf{z}(t) \rangle$$

 $a \leq t \leq b$:

$$\int_{C} F \, ds = \int_{a}^{b} F(x(t), y(t), z(t)) \, ds$$

where

$$ds = \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2} dt$$

Scalar Surface Integrals

If S is parameterized by $\mathbf{r}(u, v) = \langle x(u, v), y(u, v), z(u, v) \rangle,$ $(u, v) \in D:$

$$\iint_{S} F \, dS = \iint_{D} F |\mathbf{r}_{u} \times \mathbf{r}_{v}| \, du \, dv$$

where

$$F = F(x(u, v), y(u, v), z(u, v))$$
$$\mathbf{r}_{u} = \frac{\partial}{\partial u} \mathbf{r}(u, v)$$
$$\mathbf{r}_{v} = \frac{\partial}{\partial v} \mathbf{r}(u, v)$$

◆□▶ ◆□▶ ◆ □▶ ◆ □▶ ○ □ ○ ○ ○ ○

Vector Surface Integrals

Sneak Preview - Vector Surface Integrals

•

Vector Line Integrals

If C is parameterized by

$$\mathbf{r}(t) = \langle \mathbf{x}(t), \mathbf{y}(t), \mathbf{z}(t) \rangle,$$

 $a \leq t \leq b$:

$$\int_{C} \mathbf{F} \cdot d\mathbf{r} = \int_{a}^{b} \mathbf{F} \cdot \mathbf{T}(t) ds$$
$$\mathbf{F} = \mathbf{F}(x(t), y(t), z(t))$$
$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|}$$

Vector Surface Integrals

If S is parameterized by $\mathbf{r}(u, v) = \langle x(u, v), y(u, v), z(u, v) \rangle,$ $(u, v) \in D:$

$$\iint_{S} \mathbf{F} \cdot d\mathbf{S} = \iint_{D} \mathbf{F} \cdot \mathbf{N} \, du \, dv$$
$$\mathbf{F} = \mathbf{F}(x(u, v), y(u, v), z(u, v))$$
$$\mathbf{N} = \mathbf{r}_{u} \times \mathbf{r}_{v}$$

◆□ > ◆□ > ◆豆 > ◆豆 > ̄豆 _ のへぐ

▲ロト ▲冊 ▶ ▲ ヨ ▶ ▲ ヨ ▶ ● の Q @

Parameterized Surface Review



The parameter space D



 $\sin v \cos u \mathbf{i} + \sin v \sin u \mathbf{j} + \cos v \mathbf{k}$

A parameterized surface is traced out by $\mathbf{r}(u, v)$ where $(u, v) \in D$, a region in the uv plane.

Parameterized Surface Review



The parameter space D

A parameterized surface is traced out by $\mathbf{r}(u, v)$ where $(u, v) \in D$, a region in the uv plane.

If v is held fixed and u varies, the result is a curve along the surface.

◆□> ◆□> ◆豆> ◆豆> □豆



 $\sin v \cos u \mathbf{i} + \sin v \sin u \mathbf{j} + \cos v \mathbf{k}$

Review

Parameterized Surface Review



The parameter space D



If v is held fixed and u varies, the result is a curve along the surface.

If u is held fixed and v varies, the result is a different curve along the surface.

◆□> ◆□> ◆豆> ◆豆> □豆



 $\sin v \cos u \mathbf{i} + \sin v \sin u \mathbf{j} + \cos v \mathbf{k}$

Review

Parameterized Surface Review



The parameter space D



 $\sin v \cos u \mathbf{i} + \sin v \sin u \mathbf{j} + \cos v \mathbf{k}$

A parameterized surface is traced out by $\mathbf{r}(u, v)$ where $(u, v) \in D$, a region in the uv plane.

If v is held fixed and u varies, the result is a curve along the surface.

If u is held fixed and v varies, the result is a different curve along the surface.

Each curve has a tangent vector, so there are two independent tangent vectors

 $\mathbf{r}_u = \partial \mathbf{r} / \partial u, \quad \mathbf{r}_v = \partial \mathbf{r} / \partial v$

◆□▶ ◆□▶ ◆三▶ ◆三▶ 三三 のへぐ

Review

Parameterized Surface Review



The parameter space D



 $\sin v \cos u \mathbf{i} + \sin v \sin u \mathbf{j} + \cos v \mathbf{k}$

A parameterized surface is traced out by $\mathbf{r}(u, v)$ where $(u, v) \in D$, a region in the uv plane.

If v is held fixed and u varies, the result is a curve along the surface.

If u is held fixed and v varies, the result is a different curve along the surface.

Each curve has a tangent vector, so there are two independent tangent vectors

 $\mathbf{r}_u = \partial \mathbf{r} / \partial u, \quad \mathbf{r}_v = \partial \mathbf{r} / \partial v$

The vectors \mathbf{r}_u and \mathbf{r}_v span a tangent plane

▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ

Review

Parameterized Surface Review



The parameter space D



 $\sin v \cos u \mathbf{i} + \sin v \sin u \mathbf{j} + \cos v \mathbf{k}$

A parameterized surface is traced out by $\mathbf{r}(u, v)$ where $(u, v) \in D$, a region in the uv plane.

If v is held fixed and u varies, the result is a curve along the surface.

If u is held fixed and v varies, the result is a different curve along the surface.

Each curve has a tangent vector, so there are two independent tangent vectors

 $\mathbf{r}_u = \partial \mathbf{r} / \partial u, \quad \mathbf{r}_v = \partial \mathbf{r} / \partial v$

The vectors \mathbf{r}_u and \mathbf{r}_v span a tangent plane The normal to the tangent plane is

 $\mathbf{N} = \mathbf{r}_u \times \mathbf{r}_v$

▲ロト ▲帰ト ▲ヨト ▲ヨト 三日 - の々ぐ

Scalar Surface Integrals

If S is parameterized by $\mathbf{r}(u, v)$ for $(u, v) \in D$, and f is a function continuous in a neighborhood of S,

$$\iint_{S} f(x, y, z) \, dS = \iint_{D} F(x(u, v), y(u, v), z(u, v)) \, |\mathbf{r}_{u} \times \mathbf{r}_{v}| \, du \, dv$$

- 1. Suppose that $f(x, y, z) = g(\sqrt{x^2 + y^2 + z^2})$ where g(2) = 5. Find $\iint_S f(x, y, z) \, dS$ if *S* is the sphere $x^2 + y^2 + z^2 = 4$.
- 2. Find $\iint_S xz \, dS$ if S is the part of the plane 2x + 2y + z = 4 that lies in the first octant.
- 3. Find $\iint y^2 dS$ if S is the part of the sphere $x^2 + y^2 + z^2 = 1$ that lies above the cone $z = \sqrt{x^2 + y^2}$.

The Oriented Unit Normal to a Surface

A surface S is called an *oriented surface* if there is a unit normal vector \mathbf{n} at every point on the surface that varies continuously along the surface. Every parameterized surface has such a unit normal, given by

$$\mathbf{n} = \frac{\mathbf{r}_{u} \times \mathbf{r}_{v}}{|\mathbf{r}_{u} \times \mathbf{r}_{v}|}.$$

Every orientable surface in \mathbb{R}^3 has two possible orientations, one with n and the other with -n.



Vector Surface Integrals

Oriented Surfaces versus the Möbius Strip



The Möbius Band



August Ferdinand Möbius (1790– 1868)

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 の�?

Vector Surface Integrals

If **F** is a continuous vector field defined on an oriented surface S with unit normal vector **n**, the *surface integral of* **F** *over* S is

$$\iint_{S} \mathbf{F} \, d\mathbf{S} = \iint_{S} \mathbf{F} \cdot \mathbf{n} \, dS$$

This integral is also called the *flux* of **F** across *S*. Depending on the choice of normal, it measures either what goes *in* (inward normal) or what comes *out* (outward normal).

- 1. Find the flux of $\mathbf{F}(x, y, z) = x\mathbf{i} + y\mathbf{j} + z^2\mathbf{k}$ across S if S is the sphere of radius 1 and center at the origin
- 2. Find the flux of $\mathbf{F}(x, y, z) = y\mathbf{j} z\mathbf{k}$ across the paraboloid $y = x^2 + z^2$, $0 \le y \le 1$, and the disc $x^2 + y^2 \le 1$, y = 1