

1 Lecture 08: The squeeze theorem

- The squeeze theorem
- The limit of $\sin(x)/x$
- Related trig limits

1.1 The squeeze theorem

Example. Is the function g defined by

$$g(x) = \begin{cases} x^2 \sin(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

continuous?

Solution. If $x \neq 0$, then $\sin(1/x)$ is a composition of continuous function and thus $x^2 \sin(1/x)$ is a product of continuous function and hence continuous.

If $x = 0$, we need to have that $\lim_{x \rightarrow 0} g(x) = g(0) = 0$ in order for g to satisfy the definition of continuity. Recalling that $\sin(1/x)$ oscillates between $-1 \leq \sin(1/x) \leq 1$, we have that

$$-x^2 \leq g(x) \leq x^2$$

and since $\lim_{x \rightarrow 0} x^2 = \lim_{x \rightarrow 0} -x^2 = 0$, the theorem below tells us we have $\lim_{x \rightarrow 0} g(x) = 0$. ■

Theorem 1 (The squeeze theorem) *If f , g , and h are functions and for all x in an open interval containing c , but perhaps not at c , we have*

$$f(x) \leq g(x) \leq h(x)$$

and

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} h(x) = L,$$

then

$$\lim_{x \rightarrow c} g(x) = L.$$

We will not give a proof but it should be intuitive that if g is trapped between two functions that approach the limit L , then g also approaches that limit.

1.2 The limit of $\sin(x)/x$

We consider the limit

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x}.$$

The quotient rule for limits does not apply since the limit of the denominator is 0. Unlike our previous limits, we cannot simplify to obtain a function where we can use the direct substitution rule or another rule. Instead, we will use the squeeze theorem.


Theorem 2

$$\lim_{t \rightarrow 0} \frac{\sin(t)}{t}.$$

Proof. We start by observing that $\sin(-t)/(-t) = \sin(t)/t$, so it suffices to consider $\lim_{t \rightarrow 0^+} \sin(t)/t$.

In the figure below we draw an angle t with $0 < t < \pi/2$ and observe that we have the inequalities

$$\text{Area triangle } OAB \leq \text{Area sector } OAB \leq \text{Area triangle } OAC.$$



We have

$$\begin{aligned} \text{Area triangle } OAB &= \frac{1}{2} \sin(t) \\ \text{Area sector } OAB &= \frac{1}{2} t \\ \text{Area triangle } OAC &= \frac{1}{2} \tan(t) \end{aligned}$$

Thus we have

$$\frac{1}{2} \sin(t) \leq t/2 \leq \frac{1}{2} \tan(t).$$

Since $t > 0$, we can rearrange to obtain

$$\cos(t) \leq \frac{\sin(t)}{t} \leq 1. \quad (3)$$

and since $\sin(-t)/(-t) = \sin(t)/t$, we also have (3) if $0 < |t| < \pi/2$. Since $\lim_{t \rightarrow 0} \cos(t) = 1$, the squeeze theorem implies

$$\lim_{t \rightarrow 0} \frac{\sin(t)}{t} = 1.$$

■

1.3 Some consequences

Using this limit, we can find several related limits.

The first one will be used in the next chapter.

Example. Find the limit

$$\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x}.$$

Solution. We note that since the limit of the denominator is zero, we cannot use the quotient rule for limits. However, if we multiply and divide by $1 + \cos(x)$ and use the identity $\sin^2(x) + \cos^2(x) = 1$, we have

$$\frac{1 - \cos(x)}{x} = \frac{(1 - \cos(x))(1 + \cos(x))}{x(1 + \cos(x))} = \frac{\sin^2(x)}{x}.$$

Thus, we may use the rule for a limit of a product,

$$\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x} = \lim_{x \rightarrow 0} \frac{\sin^2(x)}{x} = \lim_{x \rightarrow 0} \sin(x) \lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 0.$$

■

Below are a few more to try

1. $\lim_{t \rightarrow 0} \frac{\sin(2t)}{t}$
2. $\lim_{t \rightarrow 0} \frac{\sin(2t)}{\sin(3t)}$
3. $\lim_{t \rightarrow 0} \frac{1 - \cos(t)}{t^2}$

February 2, 2014