

# 1 Lecture 16: Higher derivatives

## 1.1 Outline

- Definition of higher-order derivatives
- Examples

## 1.2 Higher order derivatives

The acceleration is the derivative of the derivative of position. We call this a second derivative. We often write  $f''$  for the derivative of  $f'$  and we call  $f''$  the second derivative. We can define derivatives of any order by

$$f^{(0)} = f \quad \text{and} \quad f^{(n)} = f^{(n-1)'}$$

for the result of differentiating  $n$ -times. For derivatives up to order 2 or 3, we add a prime ( $'$ ) for each derivative. But eventually we get tired of writing primes (and our readers get tired of counting) and we switch to one of the notations

$$f^{(n)} \quad \text{or} \quad \frac{d^n f}{dx^n}$$

Be careful to use the parentheses which help us to distinguish the  $n$ th-derivative  $f^{(n)}$  from the  $n$ th power  $f^n$ .

## 1.3 Examples

*Example.* Find

$$\frac{d^2}{dx^2} x^2 e^x$$

If  $f$  is defined by  $f(x) = e^x$ , find  $f^{(2014)}$ .

*Solution.* The first derivative

$$\frac{d}{dx}(x^2 e^x) = x^2 e^x + 2x e^x = (x^2 + 2x)e^x$$

Differentiating again gives

$$\frac{d}{dx}(x^2 + 2x)e^x = (2x + 2)e^x + (x^2 + 2x)e^x = (x^2 + 4x + 2)e^x.$$

After spending all day computing 2014 derivatives, we find that  $f^{(2014)}(x) = f^{(2011)}(x) = \dots = f(x) = e^x$ .

■

*Example.* Can you find a formula for

$$\frac{d^n}{dx^n} x^n, \quad \frac{d^{2n}}{dx^{2n}} x^n, \quad \frac{d^n}{dx^n} \frac{1}{x}?$$

*Solution.* Try a few examples and look for a pattern. ■

*Example.* Find a polynomial  $p(x)$  of degree 2 so that with  $f(x) = e^x$ ,  $p(0) = f(0)$ ,  $p'(0) = f'(0)$ , and  $p''(0) = f''(0)$ .

Compare the values of  $e^x$  and  $p(x)$ .

Can you suggest a better way to approximate  $e^x$ ?

*Solution.* Since  $f^{(k)}(x) = e^x$  for all  $k$ , we have  $f^{(k)}(0) = 1$  for all  $k = 0, 1, 2, \dots$ . If  $p(x) = ax^2 + bx + c$ , then we have


$$p(0) = c, \quad p'(0) = b, \quad p''(0) = 2a.$$

Since we want  $p(0) = f(0) = 1$ , we have  $c = 1$ . If  $p'(0) = 1$ , then we must have  $b = 1$  and if  $p''(0) = 1$ , we must have  $2a = 1$  or  $a = 1/2$ . Thus  $p(x) = \frac{1}{2}x^2 + x + 1$ .

Trying a few values we find

| $x$  | $e^x$          | $p(x)$  |
|------|----------------|---------|
| 0    | 1              | 1       |
| 0.01 | 1.010050167... | 1.01005 |
| 0.2  | 1.22140...     | 1.22    |
| -0.2 | 0.81873...     | 0.82    |

The graph below also indicates that the polynomial is a good approximation to the function near 0.



To do better we might look for a third degree polynomial with  $p^{(k)}(0) = f^{(k)}(0)$  for  $k = 0, 1, 2, 3$ . ■

*Example.* Find the  $n$ th derivative of  $xe^x$ . Hint: Try a few and see if you can guess a pattern.

February 20, 2014